

**CHARACTERIZATION OF CELLULASES BY
Aspergillus fumigatus USING OIL PALM FROND THROUGH
SOLID STATE FERMENTATION**

NURUL FATIHAH BINTI MOHD ZAID

**Final Year Project Report Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Science (Hons.) Biology
In the Faculty of Applied Sciences
Universiti Teknologi MARA**

JULY 2015

This Final Year Project entitled “**Characterization of Cellulases by *Aspergillus fumigatus* Using Oil Palm Frond through Solid State Fermentation**” was submitted by Nurul Fatimah binti Mohd Zaid, in partial fulfilment of the requirements for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Science, and was approved by

NoorAzrimi bin Umor
Supervisor
B. Sc. (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
72000 Kuala Pilah Negeri Sembilan

Nurul Hidayah binti Adenan
Supervisor
B. Sc. (Hons) Biology
Faculty of Applied Sciences
Universiti Teknologi MARA
72000 Kuala Pilah
Negeri Sembilan

Ilyanie binti Hj. Yaacob
Project Coordinator
B. Sc. (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
72000 Kuala Pilah Negeri Sembilan
Sembilan

Dr. Noor’Aisyah binti Abu Shah
Head of Programme
B. Sc. (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
72000 Kuala Pilah Negeri

Date: _____

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
LIST OF ABBREVIATIONS	vii
ABSTRACT	ix
ABSTRAK	x
CHAPTER 1: INTRODUCTION	
1.1 Background Study	1
1.2 Problem Statement	3
1.3 Significance of the Study	3
1.4 Objectives of the Study	4
CHAPTER 2: LITERATURE REVIEW	
2.1 Fungi	5
2.1.1 Thermophilic Fungi	5
2.1.2 Cellulolytic Fungi	7
2.2 Enzyme	7
2.3 Enzymatic Degradation of Cellulose	8
2.3.1 Cellulose	8
2.3.2 Cellulase Enzyme	9
2.4 Oil Palm	10
2.4.1 Significance of Oil Palm Biomass	11
2.4.2 Oil Palm Frond	12
2.5 Solid State Fermentation	13
CHAPTER 3: METHODOLOGY	
3.1 Materials	15
3.1.1 Microorganisms and Source	15
3.1.2 Chemicals	15
3.1.3 Apparatus	16
3.1.4 General Equipments	16

3.2	Methods	16
3.2.1	Growth Medium	17
3.2.2	Inoculum Preparation	17
3.2.3	Dry Mycelium Weight	17
3.2.4	Morphological Macroscopic and Microscopic Observations	18
3.2.5	Standard curve	18
3.2.6	Production of Thermostable Cellulase Enzyme by Using Solid State Fermentation	19
3.2.6.1	Substrate Preparation	19
3.2.6.2	Quantitative Analysis	20
3.2.6.3	Filter Paper Assay	20
3.2.6.4	CMCase assay	21
CHAPTER 4:	RESULTS AND DISCUSSION	23
4.1	Morphological Macroscopic and Microscopic Observations	25
4.2	Dry Mycelium Weight	26
4.3	Reducing Sugar Analysis	27
4.4	Production of Enzyme Assays Analysis	
CHAPTER 5:	CONCLUSIONS AND RECOMMENDATIONS	31
CITED REFERENCES		33
APPENDICES		37
CURRICULUM VITAE		43

ABSTRACT

CHARACTERIZATION OF CELLULASES BY *Aspergillus fumigatus* USING OIL PALM FROND THROUGH SOLID STATE FERMENTATION

There are lacks of microorganisms that can produce sufficient amount of three types of cellulases to break down the crystalline cellulose into glucose from the cellulosic wastes generated by agricultural and industrial activities. This study aims to characterize the thermostable enzyme produced by *Aspergillus fumigatus* (previously isolated) in solid state fermentation using oil palm frond. The thermostable cellulase enzyme activities characterized by CMCase and FPase activities were concurrently done after 3 days of fermentation. This fungi colony has grayish color and smooth walled surface through macroscopic observation whilst the structures of fungi such as vesicle, conidiophore, conidia and phialide were done by microscopic observation. CMCase assay showed higher enzyme production compared to FPase assay. The highest enzyme production for CMCase and FPase assays are 0.0535 U/mL and 0.0249 U/mL respectively. The enzyme production is low due to high temperature during fermentation and small amount of substrate has been used. The highest reducing sugar obtained is 0.865 μ mole on day 9 for reducing sugar analysis. Reducing sugar analysis showed that enzymatic hydrolysis did occurred where the cellulases broke down the cellulose into sugars. This study has proved that *Aspergillus fumigatus* can be used to degrade cellulosic wastes into soluble sugar by-products. It is suggested to prolong the enzyme production period to observe the production trends of the enzymes and increase the amount of substrate to reduce the water content to obtain higher amount of sugar produce.